

Common **WARNING SIGNS** of Mental Illness

Diagnosing mental illness isn't a straightforward science. We can't test for it the same way we can test blood sugar levels for diabetes. Each condition has its own set of unique symptoms, though symptoms often overlap. Common signs and/or symptoms can include:

- ! Feeling very sad or withdrawn for more than two weeks
- ! Trying to harm or end one's life or making plans to do so
- ! Severe, out-of-control, risk-taking behavior that causes harm to self or others
- ! Sudden overwhelming fear for no reason, sometimes with a racing heart, physical discomfort or difficulty breathing
- ! Significant weight loss or gain



- ! Seeing, hearing or believing things that aren't real*
- ! Excessive use of alcohol or drugs
- ! Drastic changes in mood, behavior, personality or sleeping habits
- ! Extreme difficulty concentrating or staying still
- ! Intense worries or fears that get in the way of daily activities

*Various communities and backgrounds might view this sign differently based on their beliefs and experiences. Some people within these communities and cultures may not interpret hearing voices as unusual.

WORRIED ABOUT YOURSELF OR SOMEONE YOU CARE ABOUT?

? If you notice any of these symptoms, it's important to ask questions



Try to understand what they're experiencing and how their daily life is impacted



Making this connection is often the first step to getting treatment

KNOWLEDGE IS POWER



Talk with a health care professional



Learn more about mental illness



Take a mental health education class



Call the NAMI HelpLine at 800-950-NAMI (6264)

50%
of all lifetime mental illness begins by age **14**

75%
by age **24**

Data from CDC, NIMH and other select sources. Find citations for this resource at nami.org/mhstats

NAMI HelpLine
800-950-NAMI (6264)

